

SPECIMEN INFORMATION

SPECIMEN:
 REQUISITION:
 LAB REF NO:

DOB:
 AGE:
 GENDER:
 FASTING:

Clinical Info:

ORDERING PHYSICIAN

CLIENT INFORMATION

DirectLabs
 4040 Florida St.
 Ste 101
 Mandeville, LA 70448

COLLECTED:
 RECEIVED:
 REPORTED:

Test Name	Result	Flag	Reference Range	Lab
PSA, POST-PROSTATECTOMY				
PSA, ICMA	0.13		ng/mL	01

REFERENCE RANGES for PSA:

LESS THAN 0.10 ng/mL AFTER RADICAL PROSTATECTOMY.

4.0 ng/mL OR LESS IN HEALTHY MALES WITHOUT PROSTATECTOMY.

PSA values obtained with different assay methods or kits cannot be used interchangeably.

This test was performed using the Beckman Coulter DxI method. PSA, ICMA is not to be used as a diagnostic procedure without confirmation of the diagnosis by another established product or procedure.

The lower limit of accurate quantification for this assay is 0.02 ng/mL. PSA values less than 0.02 ng/mL cannot be accurately measured and will be reported as less than 0.02 ng/mL. Specimens with PSA levels below the lower limit of accurate quantification should be considered as negative. In patients with a negative result for post prostatectomy PSA, serial monitoring of PSA levels at regular intervals, along with physical examinations and other tests, may help to detect recurrent prostate cancer.

Performing Laboratory Information:

01: Quest Diagnostics/Nichols SJC-San Juan Capistrano,, 33608 Ortega Hwy, San Juan Capistrano CA, phone:
 Medical Director: MD, PhD Jon Nakamoto