

SPECIMEN INFORMATION

ORDERING PHYSICIAN

SPECIMEN:

DOB:

REQUISITION:

AGE:

CLIENT INFORMATION

LAB REF NO:

GENDER:

FASTING: Unknown

DirectLabs

COLLECTED:

Clinical Info:

4040 Florida St.

RECEIVED:

Ste 101

REPORTED:

Mandeville, LA 70448

| Test Name | Result | Flag | Reference Range | Lab |
|-----------|--------|------|-----------------|-----|
|-----------|--------|------|-----------------|-----|

**YERSINIA ENTEROCOLITICA AB (IGG, IGA)**

|                      |       |  |  |    |
|----------------------|-------|--|--|----|
| Y ENTEROCOLITICA IGG | <0.90 |  |  | 03 |
| Y ENTEROCOLITICA IGA | <0.90 |  |  | 03 |

REFERENCE RANGE: <0.90

INTERPRETIVE CRITERIA:

<0.90 Antibody not detected  
 0.90 - 1.10 Equivocal  
 >1.10 Antibody detected

Yersinia enterocolitica is associated with a wide spectrum of clinical and immunological manifestations, including enteritis, colitis, reactive arthritis, and erythema nodosum. Y. enterocolitica-specific IgA and IgG are both typically detected following acute infection; IgA usually disappears after 2-3 months, whereas IgG persists for at least a year. IgA persistence is observed in some patients with chronic reactive arthritis.

This test was developed and its analytical performance characteristics have been determined by Quest Diagnostics Infectious Disease. It has not been cleared or approved by the U.S. Food and Drug Administration. The FDA has determined that such clearance or approval is not necessary. This assay has been validated pursuant to the CLIA regulations and is used for clinical purposes.

**Performing Laboratory Information:**

03: Quest Diagnostics-Infectious Disease, Inc, 33608 Ortega Highway, Building B-west Wing, San Juan Capistrano CA, phone:  
 Medical Director: MD Hollis J Batterman